Libyan Arab Jamahiriya


2010 total population: 6 355 112 Income group: Upper middle

NCD mortality*		
2008 estimates Total NCD deaths (000s)	<i>males</i> 13.5	<i>females</i> 9.6
NCD deaths under age 60 (percent of all NCD deaths)	36.3	29.3
Age-standardized death rate per 100 000		
All NCDs	743.5	525.9
Cancers	114.3	79.6
Chronic respiratory diseases	41.1	25.7
Cardiovascular diseases and diabetes	458.8	330.1

Behavioural risk factors			
2008 estimated prevalence (%)	males	females	total
Current daily tobacco smoking	45.5	0.2	23.8
Physical inactivity	35.4	53.6	44.2


Metabolic risk factors			
2008 estimated prevalence (%)	males	females	total
Raised blood pressure	45.9	39.1	42.6
Raised blood glucose	12.1	11.3	11.8
Overweight	57.8	66.2	61.9
Obesity	19.9	36.4	27.8
Raised cholesterol	33.3	33.6	33.4


Proportional mortality (% of total deaths, all ages)*





NCDs are estimated to account for 78% of all deaths.

Metabolic risk factor trends









Country capacity to address and respond to NCDs
Has a Unit / Branch / Dept in MOH with responsibility for NCDs

There is funding available for: NCD treatment and control NCD prevention and health promotion NCD surveillance, monitoring and evaluation

National health reporting system includes:

NCD cause-specific mortality Yes NCD morbidity Yes NCD risk factors Yes

Has an integrated or topic-specific policy / programme / action plan which is currently operational for:

Cardiovascular diseases	No
Cancer	DK
Chronic respiratory diseases	No
Diabetes	DK
Alcohol	Yes
Unhealthy diet / Overweight / Obesity	No
Physical inactivity	No
Tobacco	No

Number of tobacco (m)POWER measures 1/5 implemented at the highest level of achievement

Has a national, population-based cancer registry

DK = Country responded "don't know"

No

Yes

Yes

Yes

No

^{*} The mortality estimates for this country have a high degree of uncertainty because they are not based on any national NCD mortality data. The estimates are based on a combination of country life tables cause of death models, regional cause of death patterns, and WHO and UNAIDS program estimates for some major causes of death (not including NCDs).